

LCD - MoIDX: MGMT Promoter Methylation Analysis (L36192)

Links in PDF documents are not guaranteed to work. To follow a web link, please use the MCD Website.

Contractor Information

CONTRACTOR NAME	CONTRACT TYPE	CONTRACT NUMBER	JURISDICTION	STATES
Noridian Healthcare Solutions, LLC	A and B MAC	02101 - MAC A	J - F	Alaska
Noridian Healthcare Solutions, LLC	A and B MAC	02102 - MAC B	J - F	Alaska
Noridian Healthcare Solutions, LLC	A and B MAC	02201 - MAC A	J - F	Idaho
Noridian Healthcare Solutions, LLC	A and B MAC	02202 - MAC B	J - F	Idaho
Noridian Healthcare Solutions, LLC	A and B MAC	02301 - MAC A	J - F	Oregon
Noridian Healthcare Solutions, LLC	A and B MAC	02302 - MAC B	J - F	Oregon
Noridian Healthcare Solutions, LLC	A and B MAC	02401 - MAC A	J - F	Washington
Noridian Healthcare Solutions, LLC	A and B MAC	02402 - MAC B	J - F	Washington
Noridian Healthcare Solutions, LLC	A and B MAC	03101 - MAC A	J - F	Arizona
Noridian Healthcare Solutions, LLC	A and B MAC	03102 - MAC B	J - F	Arizona
Noridian Healthcare Solutions, LLC	A and B MAC	03201 - MAC A	J - F	Montana
Noridian Healthcare Solutions, LLC	A and B MAC	03202 - MAC B	J - F	Montana
Noridian Healthcare Solutions, LLC	A and B MAC	03301 - MAC A	J - F	North Dakota
Noridian Healthcare Solutions, LLC	A and B MAC	03302 - MAC B	J - F	North Dakota
Noridian Healthcare Solutions, LLC	A and B MAC	03401 - MAC A	J - F	South Dakota
Noridian Healthcare Solutions, LLC	A and B MAC	03402 - MAC B	J - F	South Dakota
Noridian Healthcare Solutions, LLC	A and B MAC	03501 - MAC A	J - F	Utah
Noridian Healthcare Solutions, LLC	A and B MAC	03502 - MAC B	J - F	Utah
Noridian Healthcare Solutions, LLC	A and B MAC	03601 - MAC A	J - F	Wyoming
Noridian Healthcare Solutions, LLC	A and B MAC	03602 - MAC B	J - F	Wyoming

LCD Information

Document Information

LCD ID

L36192

LCD Title

MoIDX: MGMT Promoter Methylation Analysis

AMA CPT / ADA CDT / AHA NUBC Copyright Statement

CPT codes, descriptions and other data only are copyright 2021 American Medical Association. All Rights Reserved. Applicable FARS/HHSARS apply.

Proposed LCD in Comment Period

N/A

Source Proposed LCD

[DL36192](#)

Original Effective Date

For services performed on or after 04/15/2016

Revision Effective Date

For services performed on or after 10/14/2021

Revision Ending Date

N/A

Retirement Date

N/A

Notice Period Start Date

02/25/2016

Notice Period End Date

04/14/2016

CMS National Coverage Policy

Title XVIII of the Social Security Act (SSA), §1862(a)(1)(A) allows coverage and payment for only those services that are considered to be reasonable and necessary for the diagnosis or treatment of illness or injury or to improve the functioning of a malformed body member.

42 CFR §410.32 Diagnostic x-ray tests, diagnostic laboratory tests, and other diagnostic tests: Conditions

CMS Internet-Only Manual, Pub. 100-02, Medicare Benefit Policy Manual, Chapter 15, §80 Requirements for Diagnostic X-Ray, Diagnostic Laboratory, and Other Diagnostic Tests, §80.1.1 Certification Changes

Coverage Guidance**Coverage Indications, Limitations, and/or Medical Necessity****Indications and Limitations of Coverage**

This policy provides limited coverage for methylation analysis for hypermethylation of the O-6-methylguanine DNA methyltransferase (MGMT) gene promoter. MGMT methylation analysis testing is considered to be reasonable and necessary for adult patients when the following criteria are met:

Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

Current Dental Terminology © 2021 American Dental Association. All rights reserved.

Copyright © 2013 - 2021, the American Hospital Association, Chicago, Illinois. Reproduced by CMS with permission. No portion of the American Hospital Association (AHA) copyrighted materials contained within this publication may be copied without the express written consent of the AHA. AHA copyrighted materials including the UB-04 codes and descriptions may not be removed, copied, or utilized within any software, product, service, solution or derivative work without the written consent of the AHA. If an entity wishes to utilize any AHA materials, please contact the AHA at 312-893-6816. Making copies or utilizing the content of the UB-04 Manual, including the codes and/or descriptions, for internal purposes, resale and/or to be used in any product or publication; creating any modified or derivative work of the UB-04 Manual and/or codes and descriptions; and/or making any commercial use of UB-04 Manual or any portion thereof, including the codes and/or descriptions, is only authorized with an express license from the American Hospital Association. To license the electronic data file of UB-04 Data Specifications, contact Tim Carlson at (312) 893-6816. You may also contact us at ub04@aha.org.

- Tumor type is high-grade malignant glioma (e.g. glioblastoma multiforme (GBM), anaplastic astrocytoma) **and**
- Patients are able to tolerate temozolomide therapy or radiation therapy, **and**
- The physician will use the MGMT testing results to decide between radiation therapy and chemotherapy alone as 1st line adjuvant treatment, or between temozolomide and other chemotherapy for 1st line adjuvant treatment

Note: This assessment is predicated on the assumption that therapy is considered beneficial for the specific patient.

Summary of Evidence

Cancer is the consequence of genetic alterations that result in a deregulation of important cellular pathways responsible for various essential functions, including cell growth, cell cycle progression, and apoptosis (programmed cell death). One result of these genetic alterations is gliomas. The treatment of high-grade gliomas, especially GBM, remains difficult as no contemporary treatments are curative. For the past several years, the standard treatment for GBM consists of maximal surgical resection, radiotherapy (RT), and concomitant and adjuvant chemotherapy with temozolomide.

Although surgical resection, RT, and chemotherapy with temozolomide are considered standard of care for most patients with high-grade glioma (including GBM and anaplastic astrocytoma), not all patients tolerate these treatments. For patients older than 70 years with a low performance rating, radiation or temozolomide alone is sometimes employed. Temozolomide treatment is not considered inferior to radiation therapy and may be tolerated better than RT by “frail” patients with low performances scores.

In patients for whom temozolomide is not the current standard of care, it has been proposed that MGMT methylation analysis can be used to predict the efficacy of temozolomide treatment. Epigenetic silencing of the MGMT DNA repair gene, by promoter methylation, leads to a lack of MGMT protein expression. Lack of MGMT protein expression immunohistochemically is related to drug responses in patients with malignant glioma treated with alkylating agents. In particular, MGMT hypermethylation is a known predictive biomarker of response to temozolomide treatment with favorable outcomes in terms of overall survival (OS) and progression free survival (PFS) in GBM patients.

MGMT promoter methylation status is a strong and independent prognostic factor in patients with newly diagnosed GBM and a clinically relevant predictive marker in the subpopulation of elderly GBM patients. MGMT promoter methylation analysis can aid in treatment decisions for patients over 70. For patients older than 70 with a good performance rating, there is evidence of benefit of temozolomide in addition to RT. In patients with lower performance, temozolomide can be used alone as it was found to be equally as effective as RT alone and it has lower toxicity for the frail population. In the temozolomide arm of both the Nordic and German trials, patients with MGMT promoter methylation had longer survival than those without. (9.7 vs 6.8 months; HR, 0.56; 95% CI, 0.34-0.93)

MGMT promoter methylation analysis also has prognostic utility. However, performing MGMT analysis is only recommended by NCCN guidelines for temozolomide guidance and not for overall prognosis prediction. Lattanzio et al⁹ confirmed that patients carrying methylation of the MGMT promoter reported a longer OS and PFS than patients with an unmethylated promoter. Wang et al²⁰ also evaluated the prognostic value of MGMT promoter methylation and TP53 mutation status found similar results.

There is still a lack of consensus on the optimal assay for reliable MGMT promoter methylation testing and a variety of tests are being used in different laboratories. According to Berghoff et al¹, pyrosequencing is the only method for which an adequately high analytical performance (high intra- and inter-laboratory repeatability and reproducibility) has been demonstrated in a fully published trial. MGMT promoter methylation testing should be performed by an experienced laboratory in which this testing has been validated.

MGMT may also be useful for determining the prognosis of colorectal cancer patients and to identify those requiring

more aggressive adjuvant therapies. Future studies will be necessary to determine its clinical utility in this area. Likewise, MGMT methylation may be an important biomarker in subsets of esophageal cancers where temozolomide may be utilized to successfully treat these patients, but where additional research on clinical utility is also needed. MGMT methylation analysis is also mentioned in the literature as a predictive marker for ovarian cancer and melanoma. However, evidence on the use of MGMT testing is unclear in these diagnoses and additional studies are needed on the clinical utility in these cancers.

Analysis of Evidence (Rationale for Determination)

Level of Evidence:

Quality - Strong

Strength - Strong

Weight - Moderate

In summary, the current literature and NCCN guidelines support the use of MGMT methylation analysis to predict the usefulness of temozolomide treatment in adult patients with high-grade gliomas.

General Information

Associated Information

Documentation Requirements

The patient's medical record must contain documentation that fully supports the medical necessity for services included within this LCD. (See "Coverage Indications, Limitations, and/or Medical Necessity") This documentation includes, but is not limited to, relevant medical history, physical examination, and results of pertinent diagnostic tests or procedures.

Documentation supporting the medical necessity should be legible, maintained in the patient's medical record, and must be made available to the MAC upon request.

Sources of Information

N/A

Bibliography

1. Berghoff AS, Hainfellner JA, Marosi C, Preusser M. Assessing MGMT methylation status and its current impact on treatment in glioblastoma. *CNS Oncol.* 2015;4(1):47-52.
2. Chang IW, Hsu CT, Lin JW, Hung CH. The prognostic impact of MGMT expression on lowgrade gangliogliomas: a clinicopathological and immunohistochemical study. *Folia Neuropathol.* 2013;51(4):275-282.
3. Furnari FB, Fenton T, Bachoo RM, et al. Malignant astrocytic glioma: genetics, biology, and paths to treatment. *Genes Dev.* 2007;21(21):2683-2710.
4. Hasina R, Surati M, Kawada I, et al. O-6-methylguanine deoxyribonucleic acid methyltransferase methylation enhances response to temozolomide treatment in esophageal cancer. *J Carcinog.* 2013;12:20.

5. Hegi ME, Diserens AC, Gorlia T, et al. MGMT gene silencing and benefit from temozolomide in glioblastoma. *N Engl J Med.* 2005;352(10):997-1003.
6. Keime-Guibert F, Chinot O, Taillandier L, et al. Radiotherapy for glioblastoma in the elderly. *N Engl J Med.* 2007; 356(15):1527-1535.
7. Krex D, Klink B, Hartmann C, et al. Long-term survival with glioblastoma multiforme. *Brain.* 2007;130(10):2596-2606.
8. Larijani L, Madjd Z, Samadikuchaksaraei A, et al. Methylation of O6-methyl guanine methyltransferase gene promoter in meningiomas comparison between tumor grades I, II, and III. *Asian Pac J Cancer Prev.* 2014;15(1):33-38.
9. Lattanzio L, Borgognone M, Mocellini C, et al. MGMT promoter methylation and glioblastoma: a comparison of analytical methods and of tumor specimens. *Int J Biol Markers.* 2015;30(2):208-216.
10. Malmstrom A, Gronberg BH, Marosi C, et al. Temozolomid versus standard 6-week radiotherapy versus hypofractionated radiotherapy in patients older than 60 years with glioblastoma: the Nordic randomised phase 3 trial. *Lancet Oncol.* 2012;13(9):916-926.
11. Martinez R, Schackert G, Yaya-Tur R, et al. Frequent hypermethylation of the DNA repair gene MGMT in long-term survivors of glioblastoma multiforme. *J Neurooncol.* 2007;83(1):91-93.
12. Mollemann M, Wolter M, Felsberg J, et al. Frequent promoter hypermethylation and low expression of the MGMT gene in oligodendroglial tumors. *Int J Cancer.* 2005;113(3):379-385.
13. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines). Central Nervous System Cancers. Version 1.2021. https://www.nccn.org/professionals/physician_gls/pdf/cns.pdf. Accessed August 25, 2021.
14. Ogino S, Kawasaki T, Brahmandam M, et al. Precision and performance characteristics of bisulfite conversion and real-time PCR (MethyLight) for quantitative DNA methylation analysis. *J Mol Diagn.* 2006;8(2):209-217.
15. Oliver JA, Ortiz R, Melguizo C, Alvarez PJ, Gomez-Millan J, Prados J. Prognostic impact of MGMT promoter methylation and MGMT and CD133 expression in colorectal adenocarcinoma. *BMC Cancer.* 2014;14:511. doi: 10.1186/1471240714511
16. Preusser M, de Ribaupierre S, Wohrer A, et al. Current concepts and management of glioblastoma. *Ann Neurol.* Jul 2011;70(1):921. doi:10.1002/ana.22425
17. Rizzo D, Scalzone M, Ruggiero A, et al. Temozolomide in the treatment of newly diagnosed diffuse brainstem glioma in children: a broken promise? *J Chemother.* 2014;27(2):106-110.
18. Roa W, Brasher PM, Bauman G, et al. Abbreviated course of radiation therapy in older patients with glioblastoma multiforme: a prospective randomized clinical trial. *J Clin Oncol.* 2004;22(9):1583-1588.
19. Sathornsumetee S, Reardon DA, Desjardins A, Quinn JA, Vredenburgh JJ, Rich JM. Molecularly targeted therapy for malignant glioma. *Cancer.* 2007;110(1):13-24.
20. Wang K, Wang YY, Ma J, et al. Prognostic value of MGMT promoter methylation and TP53 mutation in glioblastomas depends on IDH1 mutation. *Asian Pac J Cancer Prev.* 2014;15(24):10893-10898.
21. Zhang W, Lin Y, Chen B, Song SW, Jiang T. Recurrent glioblastoma of childhood treated with bevacizumab: case report and molecular features. *Childs Nerv Syst.* 2010;26(1):137-143.

Revision History Information

REVISION HISTORY DATE	REVISION HISTORY NUMBER	REVISION HISTORY EXPLANATION	REASONS FOR CHANGE
10/14/2021	R5	Under CMS National Coverage Policy added regulation CMS Internet-Only Manual, Pub. 100-02, Medicare Benefit Policy Manual, Chapter 15, §80.1.1 Certification Changes and updated descriptions to regulations.	<ul style="list-style-type: none"> • Provider Education/Guidance

REVISION HISTORY DATE	REVISION HISTORY NUMBER	REVISION HISTORY EXPLANATION	REASONS FOR CHANGE
		Under Bibliography changes were made to citations to reflect AMA citation guidelines. Formatting, punctuation, and typographical errors were corrected throughout the LCD.	
11/01/2019	R4	<p>The LCD is revised to remove CPT/HCPCS codes in the Keyword Section of the LCD.</p> <p>At this time 21st Century Cures Act will apply to new and revised LCDs that restrict coverage which requires comment and notice. This revision is not a restriction to the coverage determination; and, therefore not all the fields included on the LCD are applicable as noted in this policy.</p>	<ul style="list-style-type: none"> Other (The LCD is revised to remove CPT/HCPCS codes in the Keyword Section of the LCD.)
11/01/2019	R3	<p>11/01/2019: This LCD is being revised in order to adhere to CMS requirements per chapter 13, section 13.5.1 of the Program Integrity Manual. There has been no change in coverage with this LCD revision. Regulations regarding billing and coding were removed from the CMS National Coverage Policy section of this LCD and placed in the related Billing and Coding: MGMT Promoter Methylation Article A57433.</p>	<ul style="list-style-type: none"> Provider Education/Guidance
11/01/2019	R2	<p>As required by CR 10901, all billing and coding information has been moved to the companion article, this article is linked to the LCD.</p>	<ul style="list-style-type: none"> Revisions Due To Code Removal
01/01/2018	R1	<p>This policy is revised to comply with the 21st Century Cures Act.</p>	<ul style="list-style-type: none"> Creation of Uniform LCDs With Other MAC Jurisdiction

Associated Documents

Attachments

N/A

Related Local Coverage Documents

Articles

[A57433 - Billing and Coding: MoIDX: MGMT Promoter Methylation Analysis](#)

[A54896 - Response to Comments: MoIDX: MGMT Promoter Methylation Analysis](#)

LCDs

[DL36190 - \(MCD Archive Site\)](#)

[DL36192 - \(MCD Archive Site\)](#)

Related National Coverage Documents

N/A

Public Versions

UPDATED ON	EFFECTIVE DATES	STATUS
10/05/2021	10/14/2021 - N/A	Currently in Effect (This Version)
01/29/2020	11/01/2019 - 10/13/2021	Superseded
12/17/2019	11/01/2019 - N/A	Superseded
10/08/2019	11/01/2019 - N/A	Superseded

Some older versions have been archived. Please visit the MCD Archive Site to retrieve them.

Keywords

N/A